

# Body Condition Scoring Beef Cattle

---

Body condition scoring is a management tool that can be used to evaluate the nutritional status of beef cattle. Body condition (fat cover) is an indication of the energy reserves of a beef animal. It is important in beef production because it influences subsequent reproductive and growth performance. Cows and heifers in thin body condition at calving time are slower to rebreed, produce less colostrum, may not have sufficient nutrient reserves for maximum milk production, and are less likely to wean a live calf. Overconditioning, on the other hand, is expensive and can result in calving problems and lower dry matter intake early in lactation.

Body condition scores in beef cattle range from 1 (extremely emaciated) to 9 (very obese). Body condition can be evaluated easily by visual appraisal while driv-

ing or walking through a herd (Figure 1). It can be assessed when cattle handling may be impractical. Body condition is a more reliable indication of nutritional status than live weight. Changes in shrink, gut fill, and the weight of the fetus and fluids associated with pregnancy limit live weight from being an accurate indicator of nutritional status.

Two animals with the same body condition score may have dramatically different live weights. Similarly, cattle with the same live weight may have distinctly different body condition scores. Weight differences between condition scores vary depending on the score and where the animal is in the production cycle. These weight differences often range from 70 to 140 pounds. Percentage body fat associated with each distinct body condition score appears in **Table 1**.



Figure 1. Beef cattle producer body condition scoring a herd at pasture.



**Table 1. Beef cattle body condition score and associated body fat percent and shrunk body weight.**

Body Condition Score (BCS)	Body Fat, %	Shrunk body weight, % of BCS 5
1	3.77	77
2	7.54	81
3	11.30	87
4	15.07	93
5	18.89	100
6	22.61	108
7	26.38	118
8	30.15	130
9	33.91	144

Source: NRC, 2000. Adapted from NRC Nutrient Requirements of Beef Cattle, 7th revised edition.

Using the table above, the body weight change needed to move from one body condition score to another can be calculated. For example, a 1,200-pound cow at a body condition score of 5 would need to weigh 108 percent of its current weight to achieve a

body condition score of 6. This cow would need to gain 96 pounds ( $1,200 \times 1.08 = 1,296$ ;  $1,296 - 1,200 = 96$ ) to move up one body condition score. **Table 2** gives an example of weight differences between body condition scores for different body weights.

**Table 2. Weight changes needed to increase one beef cattle body condition score.**

Body Condition Score (BCS)	Animal Weight, pounds	Weight Change Needed to Increase 1 BCS, pounds
1	770	40
	924	48
	1,078	56
2	810	60
	972	72
	1,134	84
3	870	60
	1,044	72
	1,218	84
4	930	70
	1,116	84
	1,302	98
5	1,000	80
	1,200	96
	1,400	112
6	1,080	100
	1,296	120
	1,512	140
7	1,180	120
	1,416	144
	1,652	168
8	1,300	140
	1,560	168
	1,820	196
9	1,440	
	1,728	
	2,016	

Source: NRC, 2000. Adapted from NRC Nutrient Requirements of Beef Cattle, 7th revised edition.

Body condition depends on cattle nutritional requirements and past nutrient intake. Design nutritional programs to avoid dramatic variations in body condition scores. Recommended body condition score at calving is 5 for mature cows. Because heifers are still growing, their nutritional requirements in terms of nutrient percentages are higher than later in life.

Therefore, manage heifers to calve in a body condition score of 6. Cattle require increased percentages of total digestible nutrients, in particular, in their diets above requirements for maintenance and performance to increase body condition score. Nutrient requirements to increase body condition score of mature cows are listed in Tables 3 and 4.

**Table 3. Nutrient requirements to increase body condition score of mature beef cows from 4 to 5 during the last 90 days of pregnancy.<sup>1</sup>**

Animal Description	Dry Matter Intake (DMI)		Diet Nutrient Density					Daily Nutrients per Animal				
	DMI, lb/day	DMI, % of BW	TDN, % DM	NE <sub>m</sub> , Mcal/lb	CP, % DM	Ca, % DM	P, % DM	TDN, lb	NE <sub>m</sub> , Mcal	CP, lb	Ca, lb	P, lb
1,000	20.5	2.1	60	.59	7.7	.36	.20	12.3	12.1	1.57	.074	.040
1,100	22.0	2.0	60	.58	7.5	.35	.20	13.2	12.8	1.65	.078	.043
1,200	23.5	2.0	59	.58	7.4	.34	.19	13.9	13.6	1.74	.081	.045

<sup>1</sup>BW = total body weight = shrunk body weight or 96% full body weight, TDN = total digestible nutrients, NE<sub>m</sub> = net energy for maintenance, CP = crude protein, Ca = calcium, P = phosphorus

Source: NRC, 2000. Adapted from NRC Nutrient Requirements of Beef Cattle, 7th revised edition.

**Table 4. Nutrient requirements to increase body condition score of non-pregnant mature beef cows.<sup>1</sup>**

Animal Description	Body Condition Score (BCS)		Dry Matter Intake (DMI)		Diet Nutrient Density					Daily Nutrients per Animal				
	Current BCS	Days to gain 1 BCS	DMI, lb/day	DMI, % of BW	TDN, % DM	NE <sub>m</sub> , Mcal/lb	CP, % DM	Ca, % DM	P, % DM	TDN, lb	NE <sub>m</sub> , Mcal	CP, lb	Ca, lb	P, lb
1,000	3	30	18.8	1.9	64	.65	6.1	.28	.16	12.0	12.2	1.14	.053	.031
		60	17.7	1.8	57	.55	6.4	.23	.15	10.1	9.8	1.14	.041	.026
	4	30	20.5	2.1	66	.67	5.9	.27	.16	13.5	13.8	1.21	.056	.033
		60	19.0	1.9	58	.56	6.4	.23	.15	11.0	10.7	1.21	.044	.028
1,100	3	30	20.3	1.8	65	.66	6.0	.29	.17	13.2	13.4	1.22	.058	.034
		60	19.0	1.7	58	.56	6.4	.24	.15	11.0	10.6	1.22	.045	.029
	4	30	22.2	2.0	67	.69	5.9	.28	.16	14.9	15.3	1.30	.062	.036
		60	20.4	1.9	58	.57	6.4	.24	.15	11.8	11.6	1.30	.048	.030
1,200	3	30	21.0	1.8	65	.69	6.2	.30	.18	13.7	14.5	1.30	.063	.037
		60	20.3	1.7	58	.56	6.4	.24	.15	11.8	11.3	1.30	.048	.031
	4	30	23.5	2.0	67	.68	5.9	.28	.16	15.7	16.0	1.38	.065	.038
		60	21.8	1.8	58	.56	6.3	.23	.15	12.6	12.3	1.38	.051	.033

<sup>1</sup>BW = total body weight = shrunk body weight or 96% full body weight, TDN = total digestible nutrients, NE<sub>m</sub> = net energy for maintenance, CP = crude protein, Ca = calcium, P = phosphorus

Source: NRC, 2000. Adapted from NRC Nutrient Requirements of Beef Cattle, 7th revised edition.

## Key Places To Look for Body Condition

There are several key places to assess body condition in beef cattle. Overall body fat should be evaluated along with fat cover over the tailhead, ribs, and shoulder, and in the brisket. Muscling should be evaluated to determine if it has been broken down for energy. This occurs when cattle reach the low end of the body condition scoring scale. Visible and palpable bone structure is another essential part of body condition scoring and includes the ribs, backbone, spinous processes, transverse processes, hooks (hips), and pins.

Palpate the animal's condition over the ribs, along the backbone, and over the tailhead to assist in assigning body condition scores. Fat (condition) will be spongy to the touch. Bone structure with little or no fat cover will feel sharp to the touch. Palpation of body condition is particularly beneficial when loose hide or a thick hair coat makes visual appraisal of body condition more difficult.




Figure 2. Key places on a live beef animal to evaluate body condition.

## **Recommended Times To Body Condition Score the Herd**

Body condition scores of females in the breeding herd should fall within a range of 5 to 7 from the beginning of the calving season throughout the breeding season. Condition score cows and heifers in the herd to properly plan and adjust forage and feeding programs. This helps ensure adequate body condition for optimum reproductive performance. Following are ideal times to body condition score beef cattle:

- When calves are weaned
- 60 days prior to calving
- At calving
- At the beginning of the breeding season

Proper body condition is also important for bulls to be fertile and active breeders. Target a body condition score of 6 for bulls at the beginning of the breeding season. Monitor bull body condition during the

breeding season to identify bulls that become too thin. Hand feeding or bull rotation may be necessary to maintain adequate body condition for active breeding.

Keep good records of body condition scores in the herd on a routine basis. This is an easy and important step in developing a successful herd nutrition program. Assign feeding groups based in part on body condition score to meet cattle nutrient needs and most efficiently utilize feed and forage resources. Consider early weaning as another option for managing thin cows for successful rebreeding.

Nutrition-related costs make up a significant percentage of cash costs in most cow-calf operations in Mississippi. Monitoring body condition in the herd is a simple technique that can be used to make cost-effective herd nutrition decisions that optimize performance. For more information on body condition scoring beef cattle, contact an office of the Mississippi State University Extension Service.

### **BCS 1 = Emaciated**

No palpable fat is detectable over the spinous processes, transverse processes, ribs, or hooks. The tailhead and ribs appear very prominent.



### **BCS 2 = Poor**

Animal is still somewhat emaciated but the tailhead and ribs are less prominent. Individual spinous processes are still sharp to the touch. Some tissue cover is present over the ribs toward the top of the back.



### **BCS 3 = Thin**

Individual ribs including foreribs are easily identified but are not quite as sharp to the touch. Some fat can be felt along the spine and over the tailhead. Some tissue cover is present over the ribs toward the top of the back.



### **BCS 4 = Borderline**

Individual ribs may not be visually obvious. Individual spinous processes can be felt when palpated but feel rounded rather than sharp. Some fat cover is present over the ribs, transverse processes, and hooks.



### **BCS 5 = Moderate**

Overall appearance is generally good. Fat cover over ribs feels spongy. Palpable fat cover is present on either side of the tailhead.



### **BCS 6 = High moderate**

A high degree of palpable fat exists over the ribs and around the tailhead. Firm pressure is needed to feel the spinous processes.



**BCS 7 = Good**

Considerable fat cover is present with a fleshy overall appearance. Fat cover over the ribs and around the tailhead is very spongy. Fat "pones" or "rounds" may be starting to form along the tailhead.

**BCS 8 = Fat**

The animal is very fleshy and appears overconditioned. Palpation of the spinous processes is near impossible. Large fat deposits are present over the ribs and around the tailhead. Fat pones around the tailhead are obvious.

**BCS 9 = Extremely fat**

The overall appearance is blocky with extremely wasty and patchy fat cover. The tailhead and hooks are buried in fatty tissue with fat pones protruding. Bone structure is no longer visible and barely palpable. Large fatty deposits may even impair animal mobility.



Figure 3. Body condition scores and descriptions for beef cattle.

## References

- DeRouen, S.M., D.E. Franke, D.G. Morrison, W.E. Wyatt, D.F. Coombs, T.W. White, P.E. Humes, and B.B. Greene. 1994. Parturition body condition and weight influences on reproductive performance of first-calf beef cows. *J. Anim. Sci.* 72:1119-1125.
- Herd, D.B., and L.R. Sprott. 1986. Body condition, nutrition and reproduction of beef cows. *Texas A&M Univ. Ext. Bull.* 1526.
- Morrison, D.G., J.C. Spitzer, and J.L. Perkins. Influence of parturition body condition score change on reproduction in multiparous beef cows calving in moderate body condition. 1999. *J. Anim. Sci.* 77:1048-1054.
- National Research Council. 2000. *Nutrient Requirements of Beef Cattle*. 7th Revised Edition, 1996: Update 2000. National Academy Press. Washington, D.C.
- Tennant, C.J., J.C. Spitzer, W.C. Bridges, Jr., and J.H. Hampton. 2002. Weight necessary to change body condition scores in Angus cows. *J. Anim. Sci.* 2002. 80:2031-2035.

*The information given here is for educational purposes only. References to commercial products, trade names, or suppliers are made with the understanding that no endorsement is implied and that no discrimination against other products or suppliers is intended.*

---

*Copyright 2008 by Mississippi State University. All rights reserved. This publication may be copied and distributed without alteration for nonprofit educational purposes provided that credit is given to the Mississippi State University Extension Service.*



**MISSISSIPPI STATE**  
**UNIVERSITY**  
**EXTENSION SERVICE**

**msu**cares.com

By **Jane A. Parish**, Associate Extension/Research Professor, and **Justin D. Rhinehart**, Assistant Extension Professor, Animal and Dairy Sciences

*Discrimination based upon race, color, religion, sex, national origin, age, disability, or veteran's status is a violation of federal and state law and MSU policy and will not be tolerated. Discrimination based upon sexual orientation or group affiliation is a violation of MSU policy and will not be tolerated.*

**Publication 2508**

Extension Service of Mississippi State University, cooperating with U.S. Department of Agriculture. Published in furtherance of Acts of Congress, May 8 and June 30, 1914. MELISSA J. MIXON, Interim Director (POD-07-08)